


















# Konza Prairie Research Natural Area Soils



## Soils Legend

-  Alluvial Land
-  Benfield-Florence complex
-  Breaks-Alluvial land complex
-  Chase silty clay loam
-  Clime-Sogn complex, 5 to 20 percent slopes
-  Dwight-Irwin complex
-  Irwin silty clay loam, 4 to 8 percent slopes
-  Irwin silty clay loam, 4 to 8 percent slopes, eroded
-  Ivan silty clay loam, 1 to 3 percent slopes
-  Ivan and Kennebec silt loams
-  Reading silt loam, 0 to 1 percent slopes
-  Reading silt loam, 1 to 3 percent slopes
-  Stony steep land
-  Tully, 4 to 8 percent slopes
-  Tully, 4 to 8 percent slopes, eroded



Soil types clipped from preliminary SSURGO data. These data are not SSURGO certified. Resolution of soil type differences along county boundaries may differ from final SSURGO data. Consult "Soil Survey of Riley County and Part of Geary County, Kansas" for detailed descriptions of map unit symbols. Consult NRCS Soils Survey Center publication number 1527 for information about using SSURGO data.

Projection: UTM, zone 14, units meters, datum NAD83.

Produced by KPRNA GIS/Remote Sensing Laboratory, Div. of Biology, Kansas State University, Manhattan, KS.

Produced April 30, 1998.

